

AD-A133 102

ROTATIONAL ENERGY TRANSFER IN HIGHLY VIBRATIONALLY
EXCITED HYDROGEN FLUORIDE AND DEUTERIUM FLUORIDE(U)
WISCONSIN UNIV-MADISON DEPT OF CHEMISTRY F F CRIM
16 MAR 83 N00014-79-C-0415


1/1

UNCLASSIFIED

F/G 7/4

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

REPORT DOCUMENTATION PAGE

READ INSTRUCTIONS BEFORE COMPLETING FORM

1. REPORT NUMBER		2. GOVT ACCESSION NO.		3. RECIPIENT'S CATALOG NUMBER	
		AD-A133102			
4. TITLE (and Subtitle) ROTATIONAL ENERGY TRANSFER IN HIGHLY VIBRATIONALLY EXCITED HYDROGEN FLUORIDE AND DEUTERIUM FLUORIDE			5. TYPE OF REPORT & PERIOD COVERED Summary 6/82-5/83		
7. AUTHOR(s) F. Fleming Crim			6. PERFORMING ORG. REPORT NUMBER		
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry University of Wisconsin Madison, Wisconsin 53706			8. CONTRACT OR GRANT NUMBER(s) N00014-79-C-0415		
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Physics Program Office Arlington, Virginia 22217			10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS		
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)			12. REPORT DATE 3-16-83		
			13. NUMBER OF PAGES Five		
			15. SECURITY CLASS. (of this report)		
			15a. DECLASSIFICATION/DOWNGRADING SCHEDULE		

16. DISTRIBUTION STATEMENT (of this Report)
Approved for public release; distribution unlimited.

17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)
THE RUTH W. HOOKER
TECHNICAL LIBRARY

18. SUPPLEMENTARY NOTES
MAR 29 1983
NAVAL RESEARCH LABORATORY

19. KEY WORDS (Continue on reverse side if necessary and identify by block number)
Energy transfer, rotational relaxation, laser, hydrogen fluoride, deuterium fluoride.

20. ABSTRACT (Continue on reverse side if necessary and identify by block number)
Overtone vibration-laser double resonance measurements have provided new data on vibrational and rotational relaxation in HF(v = 2). These experiments determine the magnitudes of the total vibrational relaxation rate constants for HF(v = 1 and 2) as well as their temperature dependences. Rotational relaxation data come from these measurements as well. The temporal evolution of individual rotational states observed in the double resonance studies provides level-to-level energy transfer rate constants when analyzed using (over)

AD-A133102

DTIC FILE COPY

DTIC
SELECTED
SEP 29 1983

88 09 28

an iterative fitting scheme which incorporates scaling relations among the rate constants.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Avail and/or	
Dist	Special
A	



COLLISIONAL ENERGY TRANSFER IN HIGHLY VIBRATIONALLY EXCITED MOLECULES

(Contract No. N00014-79-C-0415)

1. Principal Investigator:

F. F. Crim
Associate Professor
Department of Chemistry
University of Wisconsin
Madison, WI 53706

2. Contract Description:

The aspect of this work funded by the Office of Naval Research is the direct measurement of rotational relaxation rates of highly vibrationally excited hydrogen fluoride and deuterium fluoride.

We expect to uncover systematic trends related to the initial rotational level of the excited molecule and to discover the influence which vibrational excitation has on rotational relaxation processes.

3. Scientific Problem:

Little experimental information is available on rotational energy transfer processes in general, and virtually none exists on the behavior of rotations in highly vibrationally excited molecules. This study will provide some of the first detailed data on rotational relaxation of molecules with large amounts of vibrational energy. As discussed below, the preparation technique and final state analysis are highly specific, so that the resulting rates represent an average over only the thermal rotational and translational energy distribution of the ground vibrational state collision partners. The rates come from a direct (time resolved) measurement of the populations of individual rotational states. Excitation of different vibrational states provides information on the influence which vibrational excitation

has on rotational relaxation rates, and produces new vibrational relaxation data, also. Our measurements will be valuable for comparison to inferential techniques, such as overtone vibration linewidth measurements, and as a test of promising theories based on modified forms of the sudden approximation. These experiments on important but as yet unexplored rotational relaxation in highly vibrationally excited molecules promise to produce unique, detailed data which extend our fundamental and practical understanding of energy transfer dynamics and test theoretical models of these processes.

4. Scientific and Technical Approach:

The detailed measurements of this study require two essential experimental capabilities: (1) preparation of a single rotational state in highly vibrationally excited molecules and (2) temporally resolved detection of the populations of other rotational levels. We prepare the desired state by direct excitation of a molecule from its ground vibrational state to an overtone vibrational level using visible photons from a pulsed dye laser or near-infrared photons produced by stimulated Raman scattering of the dye laser pulse in hydrogen gas. Because this excitation scheme relies on small transition probabilities arising from anharmonic terms in the potential and nonlinear terms in the dipole moment expansion, only a small fraction of the molecules are excited. We have demonstrated, however, that this is adequate for vibrational relaxation measurements on HF($v = 3,4,5$). (Chem. Phys. Letters 71, 258 (1980) and J. Chem. Phys. 74, 4455 (1981) and 75, 3871 (1981).) The short pulse duration (~6 ns) of our Nd:YAG/dye laser is essential to our measurements of the temporal evolution of individual rotational levels. The detection scheme employs

laser double resonance to probe a single rotational state in the vibrationally excited molecule. The exciting laser pulse and infrared probing radiation from a continuous HF(DF) laser propagate collinearly through a long (~1m) cell where transient absorption of the probe radiation mirrors the temporal evolution of an individual rotational state. After the cell, the preparation beam and probe beam are separated, and the intensity of the probe beam is recorded by a fast infrared detector and transient digitizer. A multichannel analyzer and computer accomplish the requisite signal enhancement, and analysis of the signal amplitude and form provides relaxation rates. The exciting laser selects a single vibrational-rotational level, while the probe laser interrogates an individual state also.

The specificity of excitation and detection is an essential feature of the experiment. Using this technique we can explore rotational energy transfer in HF(DF) in different vibrational levels at a variety of temperatures and with many different collision partners. These experiments provide rotational energy transfer rate constants and help to uncover the mechanisms of energy transfer. They also test theoretical descriptions of inelastic collisions in polar molecules.

5. Progress:

During the past year, we have applied our overtone vibration-laser double resonance technique to both vibrational and rotational energy transfer measurements in HF($v=2$). The vibrational relaxation studies have determined the total vibrational energy transfer rate constants for HF($v=1$ and 2) to be $k_1 = (1.46 \pm 0.1) \times 10^{-12}$ and $k_2 = (19.8 \pm 1.0) \times 10^{-12}$ $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, respectively. Most importantly, these experiments

provide the first measure of the relative importance of the vibration-to-vibration (V-V) and vibration-to-translation, rotation (V-T,R) energy transfer paths. They determine that approximately 60% of the relaxation occurs by a V-V mechanism. This value is lower than that obtained in one trajectory calculation but agrees fairly well with two others and a scaling law analysis. (R. A. Copeland, D. J. Pearson, J. M. Robinson, and F. F. Crim, J. Chem. Phys. 77, 3974 (1982).) We have also developed a heated double resonance cell and determined the temperature dependence of the rate constants for vibrational energy transfer in HF($v=1$ and 2). The rate constant for $v=2$ is unique in that it shows a markedly slower decrease with increasing temperature than $v=1,3,4$, or 5 . This seems to reflect the importance of V-V energy transfer for HF($v=2$). The V-V process apparently has a different temperature dependence than the V-T,R pathway.

The rotational relaxation measurements have produced extensive data on the time evolution of individual rotational states J' in HF($v=2$) following laser excitation of molecules into a single state J . Collisions rapidly transfer population into other levels and produce a Boltzmann rotational distributions in only a few gas kinetic collisions. Some levels can fill in only a fraction of a gas kinetic collision time. To extract individual rate constants we have used several scaling relationships in an iterative routine which fits the observed signal using level-to-level rate constants derived from the scaling relations. (R. A. Copeland and F. F. Crim, J. Chem. Phys. 78 (in press).)

6. Publications:

R. A. Copeland, D. J. Pearson, Jeanne M. Robinson and F. F. Crim,
"Laser Double Resonance Measurements of Vibrational Energy Transfer
Rates and Mechanisms in HF(v = 2)", J. Chem. Phys., 77, 3974 (1982).

R. A. Copeland and F. F. Crim, "Rotational Energy Transfer in HF(v = 2):
Experimental Measurements and Fitting Law Analysis", J. Chem. Phys.
(in press).

7. Extenuating circumstances:

none

8. No unspent funds will remain at the end of the contract period.

9. Two graduate students will have received their degrees during this
contract period.

G. M. Jursich (Ph.D.)

R. A. Copeland (Ph.D.)

10. Other support:

	<u>Amount</u>	<u>Period</u>
Air Force Office of Scientific Research	\$125,000	7/82-6/83
Army Research Office	\$199,000	8/82-7/85
Department of Energy	\$103,000	9/81-8/83
Petroleum Research Fund	\$ 30,000	9/80-8/82

FILM